Possible underestimation of the relative incidence of anaphylactic reactions to benzylisoquinoline neuromuscular blocking agents (Lagneau *et al.*) 577–578
Posterior subgluteal approach to block the sciatic nerve: the controversy of 'which came first, the chicken or the egg' is alive (Singh & Sharma) 496
Pre-emptive analgesia produced by interscalene blockade. What failed: the block or the methods? (Blumenthal *et al.*) 933–934
Prophylactic continuous intravenous ephedrine infusion for elective Caesarean section under spinal anaesthesia (Leykin & Rubulotta) 257–258
Recollection of dreams after short general anaesthesia: influence on patient anxiety and satisfaction (Knight *et al.*) 934–935
Retropharyngeal or cervicomediastinal haematomas following stellate ganglion block (Okuda *et al.*) 757–759
Sickle cell disease in pregnancy (Browne *et al.*) 75–76
Spontaneous intracranial hypotension treated with a cervical epidural blood patch (Usui *et al.*) 500–502
Subacute epidural abscess after spinal cord stimulator implantation (Arxer *et al.*) 755–757
Subcutaneous emphysema at the site of central line placement due to the haematogenous spread of *Clostridium septicum* (O'Rourke *et al.*) 162–163
Subcutaneous emphysema, pneumomediastinum and pneumothorax after laryngeal laser surgery (Sánchez-Etayo *et al.*) 753–754
Survey of antibiotic prophylaxis in cardiac surgery (Lehot *et al.*) 166–167
Temporary pacemakers for non-cardiac surgery (Csontos *et al.*) 581–582
The use of an airway exchange catheter in the management of acute airway obstruction (Dobson) 937–938
Tramadol versus meperidine in the treatment of shivering during spinal anaesthesia (Kaya *et al.*) 332–333
Two episodes of bronchospasm during anaesthesia in asthmatic patients (Yanagidate & Dohi) 579–580

Editorials

Cardiological aspects in preoperative anaesthesiological evaluation: old heroes, new shadows (Reinecke *et al.*) 595–599
Human immobilization: is the experience in Moscow just the beginning? (Stanley) 427–428
Non-heart-beating organ donation (Steven & Pace) 855–858
Resuscitative hypothermia after cardiac arrest in adults (Gwinnutt & Nolan) 511–514
SESAM – The Society in Europe for Simulation Applied to Medicine (Rall *et al.*) 763
Training, assessment and accreditation in anaesthesiology and the implications for the European Union (Simpson) 679–681

Errata

Comparison between nasal and intravenous desmopressin for the treatment of aminosalicylic acid-induced platelet dysfunction (Schulz-Stübner *et al*; 2002 **19**, 647–651) 81
COX-2-selective inhibitors: clinical relevance in surgical and acute pain (Rowbotham; 2002 **19 (Suppl. 25)**: 11–20) 995
Increasing the injection volume by dilution improves the onset of motor blockade, but not sensory blockade of ropivacaine for brachial plexus block (Krenn *et al*; 2003 **20**, 21–25) 258
Prophylactic continuous intravenous ephedrine infusion for elective Caesarean section under spinal anaesthesia (Leykin & Rubulotta; 2003 **20**, 257–258) 426
Suprascapular nerve block or a piroxicam patch for shoulder tip pain after day case laparoscopic surgery (Hong & Lee; 2003 **20**, 234–238) 426

Original Articles

A prospective, randomized, double-blind comparison of epidural levobupivacaine 0.5% with epidural ropivacaine 0.75% for lower limb procedures (Peduto *et al.*) 979–983
Adsorption of sufentanil to epidural filters and catheters (Westphal *et al.*) 124–126
Albumin administration – what is the evidence of clinical benefit? A systematic review of randomized controlled trials (Haynes *et al.*) 771–793
Anaesthesia for day case surgery: a survey of adult clinical practice in the UK (Payne *et al.*) 311–324
Anaesthesia for day case surgery: a survey of paediatric clinical practice in the UK (Payne *et al.*) 325–330
Anaesthetic and haemodynamic effects of continuous spinal versus continuous epidural anaesthesia with prilocaine (Reisli *et al.*) 26–30
Analgesic effects of intrathecal neostigmine in perianal surgery (Yegin *et al.*) 404–408
Bispectral index monitoring in patients undergoing cardiac surgery under cardiopulmonary bypass (Puri & Murthy) 451–456
Bispectral index-guided administration of anaesthesia: comparison between remifentanil/propofol and remifentanil/isoflurane (Schneider *et al.*) 624–630
Breathing pattern and workload during automatic tube compensation, pressure support and T-piece trials in weaning patients (Kuhlen *et al.*) 10–16
Cerebral vascular reactivity response to anaesthetic induction with propofol in patients with intracranial space-occupying lesions and vascular malformations (Schmieder *et al.*) 457–460
Changes in gas concentrations in the Brandt endotracheal tube cuff during and after anaesthesia with nitrous oxide (Karasawa *et al.*) 391–395
Clinical test to confirm tracheal intubation: a new method to confirm endotracheal intubation in the absence of capnography (Baigel & Safranski) 475–477
Closing the audit loop – prevention of perioperative hypothermia: audit and reaudit of perioperative hypothermia (Gallagher *et al.*) 750–752
Comparative study of the antiemetic efficacy of ondansetron, propofol and midazolam in the early postoperative period (Unlugenc *et al.*) 668–673
Comparison between the Datex-Ohmeda M-NMT® module and a force-displacement transducer for monitoring neuromuscular blockade (Motamed *et al.*) 467–469
Comparison of clevidipine with sodium nitroprusside in the control of blood pressure after coronary artery surgery (Powroznyk *et al.*) 697–703
Comparison of oral dolasetron and ondansetron in the prophylaxis of postoperative nausea and vomiting in children (Karamanlioglu *et al.*) 831–835
Comparison of the effects of sevoflurane and total intravenous anaesthesia in percutaneous nephrolithotomy (Atici & Aribogan) 653–657
Comparison of the effects of two intrathecal anaesthetic techniques for transurethral prostatectomy on haemodynamic and pulmonary function (Walsh *et al.*) 560–564
Cortical activity assessed by Narcotrend® in relation to haemodynamic responses to tracheal intubation at different stages of cortical suppression and reflex control (Raymondos *et al.*) 44–51
COX 2 selectivity of non-steroidal anti-inflammatory drugs and perioperative blood loss in hip surgery. A randomized comparison of indomethacin and meloxicam (Weber *et al.*) 963–966
Cricoid yoke: the effect of surface area and applied force on discomfort experienced by conscious volunteers (Campbell *et al.*) 52–55
Derecruitment of the lung induced by stepwise lowering of positive end-expiratory pressure in patients with adult respiratory distress syndrome (De Robertis *et al.*) 794–799

Desflurane increases heart rate independent of sympathetic activity in dogs (Picker *et al.*) 945–951
Desflurane inhibits platelet function *in vitro* similar to halothane (Berlet *et al.*) 878–883
Difficult airway management patterns among attending anaesthetists practising in Israel (Ezri *et al.*) 619–623
Disposable laryngoscope blades do not interfere with ease of intubation in scheduled general anaesthesia patients (Galinski *et al.*) 731–735
Dopamine stabilizes milrinone-induced changes in heart rate and arterial pressure during anaesthesia with isoflurane (Karasawa *et al.*) 120–123
Effect of bispectral index monitoring on sevoflurane consumption (Başar *et al.*) 396–400
Effect of catheter design on tracheal pressures during tracheal gas insufflation (Dyer *et al.*) 740–744
Effect of dexamethasone on motor brachial plexus block with bupivacaine and with bupivacaine-loaded microspheres in a sheep model (Estebe *et al.*) 305–310
Effect of sevoflurane on human neutrophil apoptosis (Tyther *et al.*) 111–115
Effect of sevoflurane on the *ex vivo* secretion of TNF-α during and after coronary artery bypass surgery (El Azab *et al.*) 380–384
Effect of the neuronal nitric oxide synthase inhibitor 7-nitroindazole on the righting reflex ED_{50} and minimum alveolar concentration during sevoflurane anaesthesia in rats (Kobayashi *et al.*) 212–219
Effects of bupivacaine used with sevoflurane on the rhythm and contractility in the isolated rat heart (Bozkurt *et al.*) 199–204
Effects of calcium and magnesium pretreatment on hyperkalaemic cardiac arrest in rats (Hollmann *et al.*) 606–611
Effects of different doses of oral ketamine for premedication of children (Turhanoğlu *et al.*) 56–60
Effects of fenoldopam on renal blood flow and its function in a canine model of rhabdomyolysis (Murray *et al.*) 711–718
Effects of nicardipine and diltiazem on the bispectral index and 95% spectral edge frequency (Hirota *et al.*) 809–812
Effects of pressure- and volume-controlled inverse ratio ventilation on haemodynamic variables, intracranial pressure and cerebral perfusion pressure in rabbits: a model of subarachnoid haemorrhage under isoflurane anaesthesia (Taplu *et al.*) 690–696
Effects of propofol on the systolic and diastolic performance of the postischaemic, reperfused myocardium in rabbits (Leather *et al.*) 191–198
Effects of repeated anaesthesia on central cholinergic function in the rat cerebral cortex (Hanning *et al.*) 93–97
Effects of rolipram, pimobendan and zaprinast on ischaemia-induced dysrhythmias and on ventricular cyclic nucleotide content in the anaesthetized rat (Carceles *et al.*) 205–211
Effects of ropivacaine on human neutrophil function: comparison with bupivacaine and lidocaine (Mikawa *et al.*) 104–110
Effects of sevoflurane on cerebral blood flow and cerebral metabolic rate of oxygen in human beings: a comparison with isoflurane (Oshima *et al.*) 543–547
Effects of xenon on mesenteric blood flow (Bogdanski *et al.*) 98–103
Efficacy of tramadol versus meperidine for pain relief and safe recovery after adenotonsillectomy (Özer *et al.*) 920–924
Elimination of alfentanil delivered by infusion is not altered by the chronic administration of atorvastatin (McDonnell *et al.*) 662–667
Epidural levobupivacaine, ropivacaine and bupivacaine in combination with sufentanil in early labour: a randomized trial (Camorcia & Capogna) 636–639
Excitatory effects of fentanyl upon the rat electroencephalogram and auditory-evoked potential responses during anaesthesia (Antunes *et al.*) 800–808
Factors determining the duration of tracheal intubation in cardiac surgery: a single-centre sequential patient audit (Naughton *et al.*) 225–233
Gastric mucosal-to-end-tidal PCO_2 difference during major abdominal surgery: influence of the arterial-to-end-tidal PCO_2 difference? (Lebuffe *et al.*) 147–152
General anaesthesia or spinal anaesthesia for outpatient urological surgery (Erhan *et al.*) 647–652
Haemodynamic response to a small intravenous bolus injection of epinephrine in cardiac surgical patients (Linton & Linton) 298–304
High-affinity block of voltage-operated rat IIA neuronal sodium channels by 2,6 di-*tert*-butylphenol, a propofol analogue (Haeseler & Leuwer) 220–224
High-dose midazolam and the attenuation of the contractile response to vasoconstrictors in coronary artery segments (Klockgether-Radke *et al.*) 289–293
Implications of the use of neuromuscular transmission monitoring on immediate postoperative extubation in off-pump coronary artery bypass surgery (Cammu *et al.*) 884–890
Improvement in postoperative pain relief by the addition of midazolam to an intrathecal injection of buprenorphine and bupivacaine (Shah *et al.*) 904–910
Increasing the injection volume by dilution improves the onset of motor blockade, but not sensory blockade of ropivacaine for brachial plexus block (Krenn *et al.*) 21–25
Induced arterial hypotension for interventional thoracic aortic stent-graft placement: impact on intracranial haemodynamics and cognitive function (von Knobelsdorff *et al.*) 134–140
Inflammation increases sufentanil requirements during surgery for inflammatory bowel diseases (Guidat *et al.*) 957–962
Inhalational induction of anaesthesia with 8% sevoflurane in children: conditions for endotracheal intubation and side-effects (Wappler *et al.*) 548–554
Injection pain of rocuronium and vecuronium is evoked by direct activation of nociceptive nerve endings (Blunk *et al.*) 245–253
Intraoperative catecholamine release in brain-dead organ donors is not suppressed by administration of fentanyl (Fitzgerald *et al.*) 952–956
Intraperitoneal application of bupivacaine plus morphine for pain relief after laparoscopic cholecystectomy (Hernández-Palazón *et al.*) 891–896
Intrathecal fentanyl added to intrathecal bupivacaine for day case surgery: a randomized study (Goel *et al.*) 294–297
Intravenous patient-controlled analgesia after thoracotomy: a comparison of morphine with tramadol (Erolçay & Yüceyar) 141–146
Levobupivacaine 0.25% compared with ropivacaine 0.25% by the caudal route in children (Astuto *et al.*) 826–830
Modelling the pharmacodynamic interaction between remifentanil and propofol by EEG-controlled dosing (Fechner *et al.*) 373–379
Multicentre evaluation of *in vitro* contracture testing with bolus administration of 4-chloro-*m*-cresol for diagnosis of malignant hyperthermia susceptibility (Wappler *et al.*) 528–536
Nerve-stimulator guided paravertebral blockade vs. general anaesthesia for breast surgery: a prospective randomized trial (Naja *et al.*) 897–903
Neural nets and prediction of the recovery rate from neuromuscular block (Santanen *et al.*) 87–92
Neuromuscular block and relative concentrations of mivacurium isomers under isoflurane versus propofol anaesthesia (Ledowski *et al.*) 821–825
Nitrous oxide concentrations in maternal and fetal blood during Caesarean section (Karasawa *et al.*) 555–559
Nitrous oxide/oxygen mixture and the prevention of pain during injection of propofol (Harmon *et al.*) 158–161
Ondansetron disintegrating tablets of 8 mg twice a day for 3 days did not reduce the incidence of nausea or vomiting after laparoscopic surgery (Thagaard *et al.*) 153–157
Opioid-induced respiratory depression is associated with increased tidal volume variability (Bouillon *et al.*) 127–133

Oral premedication with fentanyl may be a safe and effective alternative to oral midazolam (Tamura *et al.*) 482–486
Outcome for cardiothoracic surgical patients requiring multidisciplinary intensive care (Roche *et al.*) 719–725
Pain on injection of propofol: comparison of metoprolol with lidocaine (Aşık, *et al.*) 487–489
Pattern of renal dysfunction associated with myocardial revascularization surgery and cardiopulmonary bypass (Faulí *et al.*) 443–450
Perioperative headache and day case surgery (Gill *et al.*) 401–403
Perioperative intravenous adenosine infusion to extend postoperative analgesia in brachial plexus block (Apan *et al.*) 916–919
Pharmacokinetic-based total intravenous anaesthesia using remifentanil and propofol for surgical myocardial revascularization (Guarracino *et al.*) 385–390
Pharmacokinetics and pharmacodynamics of the new propofol prodrug GPI 15715 in rats (Schywalsky *et al.*) 182–190
Pharmacological modification of sodium channels from the human heart atrium in planar lipid bilayers: electrophysiological characterization of responses to batrachotoxin and pentobarbital (Wartenberg *et al.*) 354–362
Postoperative pain management with intravenous patient-controlled morphine: comparison of the effect of adding magnesium or ketamine (Ünlügenç *et al.*) 416–421
Prediction of difficult tracheal intubation (Iohom *et al.*) 31–36
Preliminary experience of combined peri- and retrobulbar block in surgery for penetrating eye injuries (Niemi-Murola *et al.*) 478–481
Preoperative oral celecoxib versus preoperative oral rofecoxib for pain relief after thyroid surgery (Karamanlıoğlu *et al.*) 490–495
Propofol versus remifentanil for monitored anaesthesia care during colonoscopy (Moerman *et al.*) 461–466
Prospective, randomized, controlled evaluation of the preventive effects of positive end-expiratory pressure on patient oxygenation during one-lung ventilation (Mascotto *et al.*) 704–710
QTc dispersion as a marker for medical complications after severe subarachnoid haemorrhage (Macmillan *et al.*) 537–542
Randomized comparison between sevoflurane anaesthesia and unilateral spinal anaesthesia in elderly patients undergoing orthopaedic surgery (Casati *et al.*) 640–646
Recollection of dreams after short general anaesthesia: influence on patient anxiety and satisfaction (Hellwagner *et al.*) 282–288
Remifentanil versus alfentanil in total intravenous anaesthesia for day case surgery (Alper *et al.*) 61–64
Remifentanil–propofol versus sufentanil–propofol anaesthesia for supratentorial craniotomy: a randomized trial (Gerlach *et al.*) 813–820
Respiratory mechanics in morbid obese patients with chronic obstructive pulmonary disease and hypertension during pneumoperitoneum (Salihoglu *et al.*) 658–661
Role of history and physical examination in preoperative evaluation (van Klei *et al.*) 612–618
Sedation for neuroradiology revisited: comparison of three techniques for cerebral angiography (Bewlay & Laurence) 726–730
Sevoflurane but not isoflurane can reduce prostacyclin production of endothelial cells (Heindl *et al.*) 116–119
Simple method to determine the size of the laryngeal mask airway in children (Gallart *et al.*) 570–574
Spinal cord stimulation does not change peripheral skin blood flow in patients with neuropathic pain (Ather *et al.*) 736–739
Stability of the LMA-ProSeal® and standard laryngeal mask airway in different head and neck positions: a randomized crossover study (Brimacombe & Keller) 65–69
Sub-Tenon's block reduces both intraoperative and postoperative analgesia requirement in vitreo-retinal surgery under general anaesthesia (Farmery *et al.*) 973–978
Sufentanil added to hyperbaric bupivacaine for subarachnoid block in Caesarean section (de Assunção Braga *et al.*) 631–635
Suprascapular nerve block or a piroxicam patch for shoulder tip pain after day case laparoscopic surgery (Hong & Lee) 234–238
Target-controlled infusion of propofol for fibreoptic intubation (Knolle *et al.*) 565–569
Testing and modelling the interaction of alfentanil and propofol on the EEG (Schwilden *et al.*) 363–372
The effect of intrathecal medetomidine on small bowel transit in the rat (Bilir *et al.*) 911–915
The efficacy of nicorandil, calcium chloride and nitroglycerin in treatment of ropivacaine-induced cardiotoxicity (Porter *et al.*) 939–944
The 'TEI-index' is preload dependent and can be measured by transoesophageal echocardiography during mechanical ventilation (Lutz *et al.*) 872–877
Time required to achieve a stable cuff pressure by repeated aspiration of the cuff during anaesthesia with nitrous oxide (Karasawa *et al.*) 470–474
Total intravenous anaesthesia with ketamine–midazolam versus halothane–nitrous oxide–oxygen anaesthesia for prolonged abdominal surgery (Shorrab & Atallah) 925–931
Tracheal intubation without muscle relaxants: remifentanil or alfentanil in combination with propofol (Erhan *et al.*) 37–43
Trazodone hydrochloride attenuates thermal hyperalgesia in a chronic constriction injury rat model (Okuda *et al.*) 409–415
Treatment of hiccup during general anaesthesia or sedation: a qualitative systematic review (Kranke *et al.*) 239–244
Trendelenburg positioning after cardiac surgery: effects on intrathoracic blood volume index and cardiac performance (Reuter *et al.*) 17–20
User acceptance of an anaesthesia information management system (Quinzio *et al.*) 967–972
What pain scales do nurses use in the postanaesthesia care unit? (Aubrun *et al.*) 745–749

Reviews

A history of neuraxial administration of local analgesics and opioids (Brill *et al.*) 682–689
Anaesthetic agents in adult day case surgery (Pollard *et al.*) 1–9
Anaesthetic implications of anticancer chemotherapy (Kvolik *et al.*) 859–871
Anaesthetic strategies to reduce perioperative blood loss in paediatric surgery (Weber *et al.*) 175–181
Basic concepts of ion channel physiology and anaesthetic drug effects (Friederich) 343–353
Fluid therapy in sepsis with capillary leakage (Marx) 429–442
Important physical principles in anaesthesiology (Manohin & Manohin) 259–281
Muscle relaxants in neurosurgical anaesthesia: a critical appraisal (Hans & Bonhomme) 600–605
Risk stratification for adverse outcome in cardiac surgery (Heijmans *et al.*) 515–527
Strategies to reduce perioperative blood loss related to non-surgical bleeding (Menzebach *et al.*) 764–770

European Journal of Anaesthesiology 2003; **20**: 1000–1008
© 2003 European Academy of Anaesthesiology
ISSN 0265-0215

Author Index

Adnet, F. *see* Galinski, M.
Ahrens, K. *see* Ledowski, T.
Akamatsu, H. *see* Mikawa, K.
Akpinar, S. *see* Başar, H.
Alagöl, A. *see* Karamanlıoğlu, B.
Albrecht, S. *see* Fechner, J.
Albrecht, S. *see* Schwilden, H.
Aldegheri, G. *see* Casati, A.
Aleixandre, F. *see* Carceles, M. D.
Alkış, N. *see* Aşık, İ.
Alper, I. *see* Erhan, E.
Alper, I., Erhan, E., Ugur, G. & Ozyar, B. Remifentanil versus alfentanil in total intravenous anaesthesia for day case surgery, 61
Altunkan, A. A. *see* Özer, Z.
Alvarez, L. *see* Faulí, A.
Anadolu, O. *see* Erhan, E.
Anbarci, O. *see* Apan, A.
Andrews, P. J. D. *see* Macmillan, C. S. A.
Anetseder, M. *see* Wappler, F.
Antunes, L. M., Roughan, J. V. & Flecknell, P. A. Excitatory effects of fentanyl upon the rat electroencephalogram and auditory-evoked potential responses during anaesthesia, 800
Anzawa, N., Hirota, K., Kitayama, M., Kushikata, T. & Matsuki, A. Fentanyl-mediated reduction in the bispectral index and 95% spectral edge frequency is age-dependent, 167
Apan, A. *see* Başar, H.
Apan, A., Ozcan, S., Buyukkocak, U., Anbarci, O. & Başar, H. Perioperative intravenous adenosine infusion to extend postoperative analgesia in brachial plexus block, 916
Aps, C. *see* Naughton, C.
Arar, C. *see* Karamanlıoğlu, B.
Arena, C. *see* Astuto, M.
Aribogan, A. *see* Atici, S.
Arkan, A. *see* Taplu, A.
Arxer, A., Busquets, C., Vilaplana, J. & Villalonga, A. Subacute epidural abscess after spinal cord stimulator implantation, 755
Aşık, İ., Alkış, N. & Elhan, A. H. A reply, 675
Aşık, I., Yörükoğlu, D., Gülay, I. & Tulunay, M. Pain on injection of propofol: comparison of metoprolol with lidocaine, 487
Astuto, M., Disma, N. & Arena, C. Levobupivacaine 0.25% compared with ropivacaine 0.25% by the caudal route in children, 826
Atallah, M. M. *see* Shorrab, A. A.
Ather, M., Di Vadi, P., Light, D., Wedley, J. R. & Hamann, W. C. Spinal cord stimulation does not change peripheral skin blood flow in patients with neuropathic pain, 736
Atici, S. & Aribogan, A. Comparison of the effects of sevoflurane and total intravenous anaesthesia in percutaneous nephrolithotomy, 653
Aubrun, F., Paqueron, X., Langeron, O., Coriat, P., Riou, B. What pain scales do nurses use in the postanaesthesia care unit?, 745
Audibert, J. *see* Kierzek, G.
Auroy, Y. *see* Villevieille, T.
Ayuso, A. *see* Sánchez-Etayo, G.

Baigel, G. & Safranski, J. Clinical test to confirm tracheal intubation: a new method to confirm endotracheal intubation in the absence of capnography, 475
Baigel, G. D. Cyproheptadine and the treatment of an unconscious patient with the serotonin syndrome, 586
Balogh, B. *see* Krenn, H.
Banzhaf, A. *see* Quinzio, L.
Barnes, N. W. *see* Picton, P.
Baroncini, S. *see* Peduto, V. A.
Barvais, L. *see* Fattorutto, M.
Başar, H. *see* Apan, A.
Başar, H., Ozcan, S., Buyukkocak, U., Akpinar, S. & Apan, A. Effect of bispectral index monitoring on sevoflurane consumption, 396
Bastien, O. *see* Lehot, J. J.
Baur, C. P. *see* Wappler, F.
Becker, B. F. *see* Heindl, B.
Benhamou, D. *see* Villevieille, T.
Benson, M. *see* Quinzio, L.
Berlet, T., Krah, A., Börner, U. & Gathof, B. S. Desflurane inhibits platelet function *in vitro* similar to halothane, 878
Bertini, L. *see* di Benedetto, P.
Bewlay, M. A. & Laurence, A. S. Sedation for neuroradiology revisited: comparison of three techniques for cerebral angiography, 726
Bhageerutty, K. *see* Tramoni, G.
Bhardwaj, N. *see* Goel, S.
Bhattacharya, A. *see* Tyagi, A.
Bilgin, E. *see* Özer, Z.
Bilir, A. *see* Bilir, L.
Bilir, L., Yelken, B., Guleç, S., Bilir, A. & Ekemen, S. The effect of intrathecal medetomidine on small bowel transit in the rat, 911
Bito, H. *see* Kobayashi, S.
Bizzarri, M. *see* Mascotto, G.
Blobner, M. *see* Bogdanski, R.
Blokland, A. *see* Hanning, C. D.
Blumenthal, S., Nadig, M. & Borgeat, A. Pre-emptive analgesia produced by interscalene blockade. What failed: the block or the methods?, 933
Blunk, J. A., Seifert, F., Schmelz, M., Reeh, P. W. & Koppert, W. Injection pain of rocuronium and vecuronium is evoked by direct activation of nociceptive nerve endings, 245
Bogar, L. *see* Csontos, Cs.
Bogdanski, R., Blobner, M., Fink, H. & Kochs, E. Effects of xenon on mesenteric blood flow, 98
Bonhomme, V. *see* Hans, P.
Booke, M. *see* Menzebach, A.
Booke, M. *see* Weber, T. P.
Booth, M. G. *see* Gallagher, G. A.
Borgeat, A. *see* Blumenthal, S.
Börner, U. *see* Berlet, T.
Bouillon, T., Bruhn, J., Roepcke, H. & Hoeft, A. Opioid-induced respiratory depression is associated with increased tidal volume variability, 127
Bozkurt, P., Süzer, Ö., Ekici, E., Demirci, Ö., Kaya, G. & Hacibekiroğlu, M. Effects of bupivacaine used with sevoflurane on the rhythm and contractility in the isolated rat heart, 199
Braga Potério, G. M. *see* de Assunção Braga, A. F.
Breithardt, G. *see* Reinecke, H.
Briggs, L. *see* Browne, I.
Brill, S., Gurman, G. M. & Fisher, A. A history of neuraxial administration of local analgesics and opioids, 682
Brimacombe, J. & Costa E. Silva, L. Pharyngeal necrosis with the laryngeal mask airway, 502

Brimacombe, J. & Keller, C. Stability of the LMA-ProSeal® and standard laryngeal mask airway in different head and neck positions: a randomized crossover study, 65
Brodner, G. *see* Westphal, M.
Browne, I., Byrne, H. & Briggs, L. Sickle cell disease in pregnancy, 75
Bruhn, J. *see* Bouillon, T.
Brüssel, T. *see* Weber, T. P.
Buerkle, H. *see* Westphal, M.
Buggy, D. J. *see* Gill, P. S.
Bullmann, V. *see* Weber, T. P.
Burak, E. *see* Schywalsky, M.
Busquets, C. *see* Arxer, A.
Buyukkocak, U. *see* Apan, A.
Buyukkocak, U. *see* Başar, H.
Byrne, H. *see* Browne, I.

Camci, E. *see* Tugrul, M.
Çamdeviren, H. *see* Özer, Z.
Cammu, G., De Keersmaecker, K., Casselman, F., Coddens, J., Hendrickx, J., Van Praet, F. & Deloof, T. Implications of the use of neuromuscular transmission monitoring on immediate postoperative extubation in off-pump coronary artery bypass surgery, 884
Camorcia, M. & Capogna, G. Epidural levobupivacaine, ropivacaine and bupivacaine in combination with sufentanil in early labour: a randomized trial, 636
Campbell, A. E., Turley, A., Wilkes, A. R. & Hall, J. E. Cricoid yoke: the effect of surface area and applied force on discomfort experienced by conscious volunteers, 52
Campistol, J. M. *see* Faulí, A.
Capogna, G. *see* Camorcia, M.
Carceles, M. D., Aleixandre, F., Fuente, T., López-Vidal, J. & Laorden, M. L. Effects of rolipram, pimobendan and zaprinast on ischaemia-induced dysrhythmias and on ventricular cyclic nucleotide content in the anaesthetized rat, 205
Carozzo, A. *see* Mascotto, G.
Casati, A. *see* di Benedetto, P.
Casati, A. *see* Mascotto, G.
Casati, A. *see* Peduto, V. A.
Casati, A., Aldegheri, G., Vinciguerra, F., Marsan, A., Fraschini, G. & Torri, G. Randomized comparison between sevoflurane anaesthesia and unilateral spinal anaesthesia in elderly patients undergoing orthopaedic surgery, 640
Casselman, F. *see* Cammu, G.
Cassens, U. *see* Menzebach, A.
Cazals, C., *see* Tramoni, G.
Celik, J. *see* Reisli, R.
Censier, K. *see* Wappler, F.
Cerchierini, E. *see* Mascotto, G.
Chauvin, M. *see* Galinski, M.
Chevanne, F. *see* Estebe, J.-P.
Clément, R. *see* Estebe, J.-P.
Coddens, J. *see* Cammu, G.
Çolak, A. *see* Karamanlıoğlu, B.
Colbert, S. A. *see* Heffernan, A. M.
Columb, M. *see* Zuokumor, P.
Combes, X. *see* Motamed, C.
Cooney, C. *see* Walsh, K. H.
Corcoran, T. *see* Murray, C.
Corda, B. *see* Lagneau, F.
Coriat, P. *see* Aubrun, F.
Costa E. Silva, L. *see* Brimacombe, J.
Costa Pereira, R. I. *see* de Assunção Braga, A. F.
Cracknell, J. *see* Kranke, P.
Cremonesi, E. *see* de Assunção Braga, A. F.
Csontos, Cs., Bogar, L., & Melczer, L. Temporary pacemakers for non-cardiac surgery, 581
Cunitz, G. *see* Schmieder, K.
Cunningham, A. J. *see* Iohom, G.
Curran, A. *see* Heffernan, A. M.

da Silva Braga, F. S. *see* de Assunção Braga, A. F.
Dahn, J., Waschke, K. F., Stuck, B. A. & Hörmann, K. Fluid shifts in anaphylaxis, 331
de Assunção Braga, A. de. F., da Silva Braga, F. S., Braga Potério, G. M., Costa Pereira, R. I., Reis, E. & Cremonesi, E. Sufentanil added to hyperbaric bupivacaine for subarachnoid block in Caesarean section, 631
De Cosmo, D. *see* Guarracino, F.
De Keersmaecker, K. *see* Cammu, G.
De Lange, J. J. *see* El Azab, S. R.
De Looze, D. A. *see* Moerman, A. T.
De Robertis, E., Servillo, G., Pezza, M., Viscidi, D. & Tufano, R. Derecruitment of the lung induced by stepwise lowering of positive end-expiratory pressure in patients with adult respiratory distress syndrome, 794
De Stefani, R. *see* Guarracino, F.
De Vos, M. M. *see* Moerman, A. T.
De Wolf, D. J. *see* Moerman, A. T.
De Wolff, M. H. *see* Leather, H. A.
Delettre, D. *see* Galinski, M.
Deloof, T. *see* Cammu, G.
Dembinski, R. *see* Kuhlen, R.
Demirci, Ö. *see* Bozkurt, P.
Demiroluk, S. *see* Salihoglu, Z.
Deusch, E. *see* Krenn, H.
di Benedetto, P., Bertini, L., Fanelli, G. & Casati, A. A reply, 497
Di Vadi, P. *see* Ather, M.
Dikmen, Y. *see* Salihoglu, Z.
Dingley, J. *see* Dyer, I. R.
Dirksen, R. *see* Weber, E. W. G.
Disma, N. *see* Astuto, M.
Dobson, G. The use of an airway exchange catheter in the management of acute airway obstruction, 937
Doetsch, S. *see* Wappler, F.
Dohi, S. *see* Yanagidate, F.
Donnelly, M. *see* O'Rourke, J.
Du Plessis, L. *see* Estebe, J.-P.
Dück, M. *see* Hoffmann, E.
Durieux, M. E. *see* Hollmann, M. W.
Durieux, M. E. *see* Weber, E. W. G.
Duthie, D. J. R. *see* Smart, J. A.
Duvaldestin, P. *see* Motamed, C.
Dyer, I. R., Esmail, M., Findlay, G., Mecklenburgh, J. S. & Dingley, J. Effect of catheter design on tracheal pressures during tracheal gas insufflation, 740

Eberhart, L. H. *see* Kranke, P.
Ecoffey, C. *see* Estebe, J.-P.
Ekemen, S. *see* Bilir, L.
Ekici, E. *see* Bozkurt, P.
El Azab, S. R., Rosseel, P. M. J., De Lange, J. J., Groeneveld, A. B. J., Van Strik, R., Van Wijk, E. M. & Scheffer, G. J. Effect of sevoflurane on the *ex vivo* secretion of TNF-α during and after coronary artery bypass surgery, 380
Elhan, A. H. *see* Aşık, İ.
Elidrissi, C. *see* Schneider, G.
Elliott, R. A. *see* Payne, K.
Elliott, R. A. *see* Pollard, B. J.
Engelhardt, M. *see* Schmieder, K.
Erbay, H., Kara, C. O., Kara, I. G. & Tomatir, E. Breathing or ventilation via a mouth mask for rhinoplasty operations in the early postanaesthetic period, 759
Erbayraktar, S. *see* Taplu, A.
Erhan, E. *see* Alper, I.

Erhan, E., Ugur, G., Alper, I., Gunusen, I. & Ozyar, B. Tracheal intubation without muscle relaxants: remifentanil or alfentanil in combination with propofol, 37
Erhan, E., Ugur, G., Anadolu, O., Saklayan, M. & Ozyar, B. General anaesthesia or spinal anaesthesia for outpatient urological surgery, 647
Erkan, N. *see* Taplu, A.
Erman, M. *see* Yegin, A.
Ermert, T. *see* Westphal, M.
Erolçay, H. & Yüceyar, L. Intravenous patient-controlled analgesia after thoracotomy: a comparison of morphine with tramadol, 141
Esmail, M. *see* Dyer, I. R.
Estebe, J.-P., Le Corre, P., Clément, R., Du Plessis, L., Chevanne, F., Le Verge, R. & Ecoffey, C. Effect of dexamethasone on motor brachial plexus block with bupivacaine and with bupivacaine-loaded microspheres in a sheep model, 305
Ezri, T., Konichezky, S., Geva, D., Warters, R. D., Szmuk, P. & Hagberg, C. Difficult airway management patterns among attending anaesthetists practising in Israel, 619

Fahy, C. *see* O'Rourke, J.
Fanelli, G. *see* di Benedetto, P.
Farmery, A. D. *see* Roche, R. J.
Farmery, A. D., Shlugman, D., Rahman, R. & Rosen, P. Sub-Tenon's block reduces both intraoperative and postoperative analgesia requirement in vitreo-retinal surgery under general anaesthesia, 973
Fattorutto, M. Central neuraxial blocks and anticoagulation, 254
Fattorutto, M., Pradier, O., Jansens, J. L., Ickx, B. & Barvais, L. Plateletpheresis the day before cardiac surgery and the impairment of platelet function, 338
Faulí, A., Gomar, C., Campistol, J. M., Alvarez, L., Manig, A. M. & Matute, P. Pattern of renal dysfunction associated with myocardial revascularization surgery and cardiopulmonary bypass, 443
Fechner, J., Hering, W., Ihmsen, H., Palmaers, T., Schüttler, J. & Albrecht, S. Modelling the pharmacodynamic interaction between remifentanil and propofol by EEG-controlled dosing, 373
Fechner, J. *see* Schwilden, H.
Fechner, J. *see* Schywalsky, M.
Felbinger, T. W. *see* Reuter, D. A.
Felleiter, P. *see* Wappler, F.
Feneck, R. O. *see* Naughton, C.
Feneck, R. O. *see* Powroznyk, A. V. V.
Fernandez-Galinski, S. *see* Gallart, L.
Fiege, M. *see* Wappler, F.
Findlay, G. *see* Dyer, I. R.
Fink, H. *see* Bogdanski, R.
Fisher, A. *see* Brill, S.
Fitzgerald, R. D. *see* Krenn, H.
Fitzgerald, R. D., Hieber, C., Schweitzer, E., Luo, A., Oczenski, W. & Lackner, F. X. Intraoperative catecholamine release in brain-dead organ donors is not suppressed by administration of fentanyl, 952
Flecknell, P. A. *see* Antunes, L. M.
Foubert, L. A. *see* Moerman, A. T.
Fraschini, G. *see* Casati, A.
Frerichs, A. *see* Klockgether-Radke, A. P.
Fricker, R. *see* Wappler, F.
Friederich, P. Basic concepts of ion channel physiology and anaesthetic drug effects, 343
Frietsch, T. A reply, 423
Frings, D. P. *see* Wappler, F.
Fuente, T. *see* Carceles, M. D.
Fujii, Y. Effects of diltiazem compared with nicardipine on diaphragmatic fatigability *in vivo*, 575
Fukuda, I. *see* Karasawa, F.
Galinski, M., Adnet, F., Tran, D., Karyo, Z., Quintard, H., Delettre, D., Lebail, E., Guignard, B., Lebrault, C. & Chauvin, M. Disposable laryngoscope blades do not interfere with ease of intubation in scheduled general anaesthesia patients, 731
Gallagher, G. A., McLintock, T. & Booth, M. G. Closing the audit loop – prevention of perioperative hypothermia: audit and reaudit of perioperative hypothermia, 750
Gallart, L., Mases, A., Martinez, J., Montes, A., Fernandez-Galinski, S. & Puig, M. M. Simple method to determine the size of the laryngeal mask airway in children, 570
Garrard, C. S. *see* Roche, R. J.
Gaszyński, T., Gaszyński, W. & Strzelczyk, J. General anaesthesia with remifentanil and cisatracurium for a superobese patient, 77
Gaszyński, W. *see* Gaszyński, T.
Gathof, B. S. *see* Berlet, T.
Geldner, G. *see* Ledowski, T.
Gemlik, I. *see* Karamanlıoğlu, B.
Gerlach, K., Uhlig, T., Hüppe, M., Nowak, G., Schmitz, A., Saager, L., Grasteit, A. & Schmucker, P. Remifentanil–propofol versus sufentanil–propofol anaesthesia for supratentorial craniotomy: a randomized trial, 813
Geva, D. *see* Ezri, T.
Giebler, R. *see* Lutz, J. T.
Gill, P. S., Guest, C., Rabey, P. G. & Buggy, D. J. Perioperative headache and day case surgery, 401
Giménez-Viudes, J. *see* Hernández-Palazón, J.
Glavas-Obrovac, L. *see* Kvolik, S.
Glavin, R. *see* Rall, M.
Goel, S., Bhardwaj, N. & Grover, V. K. Intrathecal fentanyl added to intrathecal bupivacaine for day case surgery: a randomized study, 294
Goetz, A. E. *see* Reuter, D. A.
Gökmen, N. *see* Taplu, A.
Gomar, C. *see* Faulí, A.
Goodchild, C. S. *see* Shah, F. R.
Görür, K. *see* Özer, Z.
Gosavi, C. P. *see* Poddar, R. K.
Goto, F. *see* Ogawa-Okamoto, C.
Goto, F. *see* Usui, T.
Gottwald, B. *see* Quinzio, L.
Grasteit, A. *see* Gerlach, K.
Greher, M. *see* Hellwagner, K.
Greim, C.-A. *see* Kranke, P.
Grobbee, D. E. *see* van Klei, W. A.
Groeneveld, A. B. J. *see* El Azab, S. R.
Große Hartlage, M. A. *see* Weber, T. P.
Grouven, U. *see* Raymondos, K.
Grover, V. K. *see* Goel, S.
Guarracino, F., Penzo, D., De Cosmo, D., Vardanega, A. & De Stefani, R. Pharmacokinetic-based total intravenous anaesthesia using remifentanil and propofol for surgical myocardial revascularization, 385
Guest, C. *see* Gill, P. S.
Guidat, A., Fleyfel, M., Vallet, B., Desreumaux, P., Levron, J. C., Gambiez, L., Colombel, J. F. & Scherpereel, P. Inflammation increases sufentanil requirements during surgery for inflammatory bowel diseases, 957
Guignard, B. *see* Galinski, M.
Gülay, I. *see* Aşık, I.
Guleç, S. *see* Bilir, L.
Güler, T. *see* Ünlügenç, H.
Gunes, Y. *see* Unlugenc, H.
Gunusen, I. *see* Erhan, E.
Gurman, G. M. *see* Brill, S.
Gustorff, B. *see* Hellwagner, K.
Gustorff, B. *see* Knolle, E.
Gwinnutt, C. L. & Nolan, J. P. Resuscitative hypothermia after cardiac arrest in adults, 511

Haasio, J. *see* Santanen, O. A. P.
Hacibekiroğlu, M. *see* Bozkurt, P.
Haeseler, G. & Leuwer, M. High-affinity block of voltage-operated rat IIA neuronal sodium channels by 2,6 di-*tert*-butylphenol, a propofol analogue, 220
Hagberg, C. *see* Ezri, T.
Halbe, A. R. *see* Shah, F. R.
Hall, J. E. *see* Campbell, A. E.
Halsall, P. J. *see* Wappler, F.
Haltiavaara, K. M. A reply 934
Haltiavaara, K. M., Laitinen, J. O., Kaukinen, S., Viljakka, T. J., Laippala, P. J. & Luukkaala, T. H. Failure of interscalene brachial plexus blockade to produce pre-emptive analgesia after shoulder surgery, 72
Hamachi, T. *see* Karasawa, F.
Hamann, W. C. *see* Ather, M.
Hanning, C. D., Blokland, A., Johnson, M., Perry, E. K. Effects of repeated anaesthesia on central cholinergic function in the rat cerebral cortex, 93
Hans, P. & Bonhomme, V. Muscle relaxants in neurosurgical anaesthesia: a critical appraisal, 600
Harada, C., Sakuragi, T., Katori, K. & Higa, K. Impossibility of injection through an epidural catheter caused by an incorrect connection of catheter and connector, 585
Harders, A. *see* Schmieder, K.
Hardman, J. G. *see* Knight, D. J. W.
Harmon, D., Rozario, C. & Lowe, D. Nitrous oxide/oxygen mixture and the prevention of pain during injection of propofol, 158
Hartmann, B. *see* Quinzio, L.
Hartung, E. *see* Wappler, F.
Haynes, G. R., Navickis, R. J. & Wilkes, M. M. Albumin administration – what is the evidence of clinical benefit? A systematic review of randomized controlled trials, 771
Heffernan, A. M., White, M., Curran, A., Colbert, S. A. Laryngeal mask airway severed by biting, 74
Heffron, J. J. A. *see* Wappler, F.
Heijmans, J. H., Maessen, J. G. & Roekaerts, P. M. H. J. Risk stratification for adverse outcome in cardiac surgery, 515
Heindel, W. *see* Weber, T. P.
Heindl, B., Reichle, F. & Becker, B. F. Sevoflurane but not isoflurane can reduce prostacyclin production of endothelial cells, 116
Hellige, G. *see* Klockgether-Radke, A. P.
Hellwagner, K. *see* Knolle, E.
Hellwagner, K., Holzer, A., Gustorff, B., Schroegendorfer, K., Greher, M., Weindlmayr-Goettel, M., Saletu, B. & Lackner, F. X. Recollection of dreams after short general anaesthesia: influence on patient anxiety and satisfaction, 282
Hellwagner, K., Holzer, A., Gustorff, B., Schroegendorfer, K., Greher, M., Weindlmayr-Goettel, M. Saletu, B. & Lackner, F. X. A reply, 936
Helou, S. *see* Lehot, J. J.
Hempelmann, G. *see* Quinzio, L.
Henderson, K. *see* White, S.
Hendrickx, J. *see* Cammu, G.
Hennis, P. J. *see* van Klei, W. A.
Hering, W. *see* Fechner, J.
Hering, W. *see* Schwilden, H.
Hernández-Palazón, J., Tortosa, J. A., Nuño de la Rosa, V., Giménez-Viudes, J., Ramírez, G. & Robles, R. Intraperitoneal application of bupivacaine plus morphine for pain relief after laparoscopic cholecystectomy, 891
Herregods, L. L. *see* Moerman, A. T.
Heytens, L. *see* Wappler, F.
Hieber, C. *see* Fitzgerald, R. D.
Higa, K. *see* Harada, C.
Hirota, K. *see* Anzawa, N.
Hirota, K., Kabara, S., Kushikata, T., Kitayama, M., Ishihara, H. & Matsuki, A. Effects of nicardipine and diltiazem on the bispectral index and 95% spectral edge frequency, 809
Hoeft, A. *see* Bouillon, T.
Hoffmann, E., Dück, M., Oberthür, A. & Paul, M. Anaesthesia for repair of cleft lip, maxilla and palate in Nepal, 677
Hohage, H. *see* Westphal, M.
Hollmann, M. W., Strümper, D., Salmons, V. A., Washington, J. M. & Durieux, M. E. Effects of calcium and magnesium pretreatment on hyperkalaemic cardiac arrest in rats, 606
Holloway, J. *see* Powroznyk, A. V. V.
Holzer, A. *see* Hellwagner, K.
Hong, J. Y. & Lee, I. H. Suprascapular nerve block or a piroxicam patch for shoulder tip pain after day case laparoscopic surgery, 234
Hopkins, P. M. *see* Wappler, F.
Höppner, R. M. *see* von Knobelsdorff, G.
Hörmann, K. *see* Dahn, J.
Hunt, T. *see* Naughton, C.
Hunter, D. *see* Naughton, C.
Hüppe, M. *see* Gerlach, K.

Ickx, B. *see* Fattorutto, M.
Ihmsen, H. *see* Fechner, J.
Ihmsen, H. *see* Schwilden, H.
Ihmsen, H. *see* Schywalsky, M.
Ikemoto, T. *see* Karasawa, F.
Immonen, I. *see* Niemi-Murola, L.
Iohom, G. *see* Walsh, K. H.
Iohom, G., Ronayne, M. & Cunningham, A. J. Prediction of difficult tracheal intubation, 31
Ishihara, H. *see* Hirota, K.
Işik, G. *see* Ünlügenç, H.
Iwamoto, T. *see* Kobayashi, S.

Jansens, J. L. *see* Fattorutto, M.
Jellinek, H. *see* Krenn, H.
Johansson, T. *see* Lisander, B.
Johnson, M. *see* Hanning, C. D.
Jolin-Mellgård, Å. *see* Powroznyk, A. V. V.
Jost, A. *see* Quinzio, L.
Junger, A. *see* Quinzio, L.
Jürgens, E. *see* Kuhlen, R.

Kabara, S. *see* Hirota, K.
Kadiogullari, D. N. *see* Kaya, M.
Kalkman, C. J. *see* van Klei, W. A.
Kallio, H. *see* Niemi-Murola, L.
Kara, C. O. *see* Erbay, H.
Kara, I. G. *see* Erbay, H.
Karadibak, K. *see* Taplu, A.
Karakus, D. *see* Kaya, M.
Karamanlıoğlu, B. *see* Turan, A.
Karamanlıoğlu, B., Turan, A., Memiş, D. & Süt, N. Comparison of oral dolasetron and ondansetron in the prophylaxis of postoperative nausea and vomiting in children, 831
Karamanlıoğlu, B., Arar, C., Alagöl, A., Çolak, A., Gemlik, I. & Süt, N. Preoperative oral celecoxib versus preoperative oral rofecoxib for pain relief after thyroid surgery, 490
Kararmaz, A. *see* Turhanoğlu, S.
Karasawa, F. *see* Oshima, T.
Karasawa, F., Hamachi, T., Takamatsu, I. & Oshima, T. Time required to achieve a stable cuff pressure by repeated aspiration of the cuff during anaesthesia with nitrous oxide, 470
Karasawa, F., Okuda, T., Tsutsui, M., Matsuoka, N., Yamada, S., Kawatani, Y. & Satoh, T. Dopamine stabilizes milrinone-induced changes in heart rate and arterial pressure during anaesthesia with isoflurane, 120

Karasawa, F., Takita, A., Fukuda, I. & Kawatani, Y. Nitrous oxide concentrations in maternal and fetal blood during Caesarean section, 555
Karasawa, F., Takita, A., Kodama, M., Takahashi, T., Wada, H., Ikemoto, T. & Satoh, T. Changes in gas concentrations in the Brandt endotracheal tube cuff during and after anaesthesia with nitrous oxide, 391
Karner, I. *see* Kvolik, S.
Karsli, B. *see* Yegin, A.
Karyo, Z. *see* Galinski, M.
Katoh, T. *see* Kobayashi, S.
Katori, K. *see* Harada, C.
Kaukinen, S. *see* Haltiavaara, K. M.
Kawatani, Y. *see* Karasawa, F.
Kaya, G. *see* Bozkurt, P.
Kaya, M., Sariyildiz, O., Karakus, D., Özalp, G. & Kadiogullari, D. N. Tramadol versus meperidine in the treatment of shivering during spinal anaesthesia, 332
Kaya, S. *see* Turhanoğlu, S.
Kehrel, B. E. *see* Weber, T. P.
Keller, C. *see* Brimacombe, J.
Kettler, D. *see* Klockgether-Radke, A. P.
Kierzek, G., Audibert, J. & Pourriat, J.-L. Anaphylaxis after rocuronium, 169
Kilger, E. *see* Reuter, D. A.
Kirov, K. *see* Motamed, C.
Kitagawa, S. *see* Tamura, M.
Kitajima, T. *see* Okuda, Y.
Kitamura, R. *see* Tamura, M.
Kitayama, M. *see* Anzawa, N.
Kitayama, M. *see* Hirota, K.
Klingler, W. *see* Wappler, F.
Klockgether-Radke, A. P., Thudium, A., Frerichs, A., Kettler, D. & Hellige, G. High-dose midazolam and the attenuation of the contractile response to vasoconstrictors in coronary artery segments, 289
Knape, J. T. A. *see* van Klei, W. A.
Knight, D. J. W. & Hardman, J. G. Recollection of dreams after short general anaesthesia: influence on patient anxiety and satisfaction, 934
Knight, D. J. W., Moppett, I. K. & Hardman, J. G. Breathing pattern and workload during automatic tube compensation, pressure support and T-piece trials in weaning patients, 932
Knolle, E., Oehmke, M. J., Gustorff, B., Hellwagner, K. & Kress, H. G. Target-controlled infusion of propofol for fibreoptic intubation, 565
Kobayashi, S., Katoh, T., Iwamoto, T., Bito, H. & Sato, S. Effect of the neuronal nitric oxide synthase inhibitor 7-nitroindazole on the righting reflex ED_{50} and minimum alveolar concentration during sevoflurane anaesthesia in rats, 212
Koch, C. *see* Wappler, F.
Kochs, E. *see* Bogdanski, R.
Kodama, M. *see* Karasawa, F.
Konichezky, S. *see* Ezri, T.
Koppert, W. *see* Blunk, J. A.
Krah, A. *see* Berlet, T.
Kranke, P., Eberhart, L. H., Morin, A. M., Cracknell, J., Greim, C.-A. & Roewer, N. Treatment of hiccup during general anaesthesia or sedation: a qualitative systematic review, 239
Krauss, T. *see* Raymondos, K.
Krenn, H., Deusch, E., Balogh, B., Jellinek, H., Oczenski, W., Plainer-Zöchling, E. & Fitzgerald, R. D. Increasing the injection volume by dilution improves the onset of motor blockade, but not sensory blockade of ropivacaine for brachial plexus block, 21
Kress, H. G. *see* Knolle, E.
Kress, H.-G. *see* Ledowski, T.
Kuhlen, R., Max, M., Dembinski, R., Terbeck, S., Jürgens, E. & Rossaint, R. Breathing pattern and workload during automatic tube compensation, pressure support and T-piece trials in weaning patients, 10
Kuhlen, R., Max, M., Dembinski, R., Terbeck, S., Jürgens, E. & Rossaint, R. A reply, 933
Kushikata, T. *see* Anzawa, N.
Kushikata, T. *see* Hirota, K.
Kvolik, S., Glavas-Obrovac, L., Sakic, K., Margaretic, D. & Karner, I. Anaesthetic implications of anticancer chemotherapy, 859

Lackner, F. X. *see* Fitzgerald, R. D.
Lackner, F. X. *see* Hellwagner, K.
Lagneau, F., Corda, B. & Marty, J. Possible underestimation of the relative incidence of anaphylactic reactions to benzylisoquinoline neuromuscular blocking agents, 577
Laippala, P. J. *see* Haltiavaara, K. M.
Laitinen, J. O. *see* Haltiavaara, K. M.
Lamm, P. *see* Reuter, D. A.
Langeron, O. *see* Aubrun, F.
Laorden, M. L. *see* Carceles, M. D.
Latimer, R. D. *see* Powroznyk, A. V. V.
Laurence, A. S. *see* Bewlay, M. A.
Le Corre, P. *see* Estebe, J.-P.
Le Verge, R. *see* Estebe, J.-P.
Leary, T. S. *see* Sturgess, J. E.
Leather, H. A., De Wolff, M. H. & Wouters, P. F. Effects of propofol on the systolic and diastolic performance of the postischaemic, reperfused myocardium in rabbits, 191
Lebail, E. *see* Galinski, M.
Lebrault, C. *see* Galinski, M.
Lebuffe, G., Onimus, T. & Vallet, B. Gastric mucosal-to-end-tidal PCO_2 difference during major abdominal surgery: influence of the arterial-to-end-tidal PCO_2 difference?, 147
Ledowski, T., Wulf, H., Ahrens, K., Weindlmayr-Goettel, M., Kress, H.-G., Geldner, G. & Scholz, J. Neuromuscular block and relative concentrations of mivacurium isomers under isoflurane versus propofol anaesthesia, 821
Lee, I. H. *see* Hong, J. Y.
Lehmann-Horn, F. *see* Wappler, F.
Lehot, J. J., Helou, S. & Bastien, O. Survey of antibiotic prophylaxis in cardiac surgery, 166
Leuwer, M. *see* Haeseler, G.
Leykin, Y. & Rubulotta, F. Prophylactic continuous intravenous ephedrine infusion for elective Caesarean section under spinal anaesthesia, 257
Light, D. *see* Ather, M.
Linton, N. W. F. & Linton, R. A. F. Haemodynamic response to a small intravenous bolus injection of epinephrine in cardiac surgical patients, 298
Linton, R. A. F. *see* Linton, N. W. F.
Lisander, B. & Johansson, T. Do we know all about hypothermia and are studies on the subject unethical?, 851
Lönnqvist, P. A. *see* Naja, M. Z.
López-Vidal, J. *see* Carceles, M. D.
Lowe, D. *see* Harmon, D.
Luis, M. *see* Sánchez-Etayo, G.
Luo, A. *see* Fitzgerald, R. D.
Lutz, J. T., Giebler, R. & Peters, J. The 'TEI-index' is preload dependent and can be measured by transoesophageal echocardiography during mechanical ventilation, 872
Luukkaala, T. H. *see* Haltiavaara, K. M.

Macmillan, C. S. A., Andrews, P. J. D. & Struthers, A. D. QTc dispersion as a marker for medical complications after severe subarachnoid haemorrhage, 537
Maessen, J. G. *see* Heijmans, J. H.
Malkan, D. *see* McDonnell, C. G.
Manig, A. M. *see* Faulí, A.
Mann, V. *see* Wappler, F.

Manohin, A. & Manohin, M. Important physical principles in anaesthesiology, 259
Manohin, M. *see* Manohin, A.
Margaretic, D. *see* Kvolik, S.
Markos, F. *see* Murray, C.
Markos, F. *see* Porter, J. M.
Marsan, A. *see* Casati, A.
Martinez, J. *see* Gallart, L.
Marty, J. *see* Lagneau, F.
Marx, G. Fluid therapy in sepsis with capillary leakage, 429
Mascotto, G., Bizzarri, M., Messina, M., Cerchierini, E., Torri, G., Carozzo, A. & Casati, A. Prospective, randomized, controlled evaluation of the preventive effects of positive end-expiratory pressure on patient oxygenation during one-lung ventilation, 704
Mases, A. *see* Gallart, L.
Mather, S. *see* Rall, M.
Matsuki, A. *see* Anzawa, N.
Matsuki, A. *see* Hirota, K.
Matsuoka, N. *see* Karasawa, F.
Matute, P. *see* Faulí, A.
Maunuksela, E.-L. *see* Niemi-Murola, L.
Max, M. *see* Kuhlen, R.
McAdoo, J. *see* Walsh, K. H.
McDonnell, C. G., Malkan, D., Van Pelt, F. D. & Shorten, G. D. Elimination of alfentanil delivered by infusion is not altered by the chronic administration of atorvastatin, 662
McHugh, G. A. *see* Payne, K.
McLintock, T. *see* Gallagher, G. A.
Mecklenburgh, J. S. *see* Dyer, I. R.
Meeting of the Society in Europe for Simulation Applied to Medicine (SESAM), 10–11 May 2002: Santander, Spain, 836
Meier-Hellmann, A. *see* Sakka, S. G.
Meißner, A. *see* Weber, T. P.
Melczer, L. *see* Csontos, Cs.
Memiş, D. *see* Karamanlıoğlu, B.
Memiş, D. *see* Turan, A.
Mentzelopoulos, S. D., Rellos, K., Tzoufi, M. J., Michalopoulos, A. S. & Papageorgiou, E. P. A double-lumen tube technique for selective lobar isolation, 986
Menzebach, A., Cassens, U., Van Aken, H. & Booke, M. Strategies to reduce perioperative blood loss related to non-surgical bleeding, 764
Mercier, F. *see* Villevieille, T.
Messina, M. *see* Mascotto, G.
Michalopoulos, A. S. *see* Mentzelopoulos, S. D.
Mikawa, K., Akamatsu, H., Nishina, K., Shiga, M., Obara, H. & Niwa, Y. Effects of ropivacaine on human neutrophil function: comparison with bupivacaine and lidocaine, 104
Misso, S. L. *see* Powroznyk, A. V. V.
Moerman, A. T., Foubert, L. A., Herregods, L. L., Struys, M. M. R. F., De Wolf, D. J., De Looze, D. A., De Vos, M. M. & Mortier, E. P. Propofol versus remifentanil for monitored anaesthesia care during colonoscopy, 461
Moerstedt, K. *see* Reuter, D. A.
Mönk, S. *see* Rall, M.
Montanini, S. *see* Peduto, V. A.
Montes, A. *see* Gallart, L.
Moons, K. G. M. *see* van Klei, W. A.
Moore, E. W. *see* Payne, K.
Moore, E. W. *see* Pollard, B. J.
Moore, J. K. *see* Payne, K.
Moppett, I. K. *see* Knight, D. J. W.
Mori, N. *see* Tamura, M.
Morin, A. M. *see* Kranke, P.
Mortier, E. P. *see* Moerman, A. T.
Motamed, C., Kirov, K., Combes, X. & Duvaldestin, P. Comparison between the Datex-Ohmeda M-NMT® module and a force-displacement transducer for monitoring neuromuscular blockade, 467
Münte, S. *see* Raymondos, K.
Murphy, C. *see* Walsh, K. H.
Murray, C., Markos, F., Snow, H. M., Corcoran, T., Parfrey, N. & Shorten, G. D. Effects of fenoldopam on renal blood flow and its function in a canine model of rhabdomyolysis, 711
Murthy, S. S. *see* Puri, G. D.

Nadig, M. *see* Blumenthal, S.
Naja, M. Z., Ziade, M. F., Lönnqvist, P. A. Nerve-stimulator guided paravertebral blockade vs. general anaesthesia for breast surgery: a prospective randomized trial, 897
Nakamura, K. *see* Tamura, M.
Naughton, C. *see* Powroznyk, A. V. V.
Naughton, C., Reilly, N., Powroznyk, A., Aps, C., Hunt, T., Hunter, D., Parsons, R. S., Sherry, E., Spackman, D., Wielogorski, A. & Feneck, R. O. Factors determining the duration of tracheal intubation in cardiac surgery: a single-centre sequential patient audit, 225
Navickis, R. J. *see* Haynes, G. R.
Niemi-Murola, L., Immonen, I., Kallio, H. & Maunuksela, E.-L. Preliminary experience of combined peri- and retrobulbar block in surgery for penetrating eye injuries, 478
Nienaber, C. A. *see* von Knobelsdorff, G.
Nishihara, F. *see* Ogawa-Okamoto, C.
Nishina, K. *see* Mikawa, K.
Nivoche, Y. *see* Wappler, F.
Niwa, Y. *see* Mikawa, K.
Nolan, J. P. *see* Gwinnutt, C. L.
Nordlander, M. *see* Powroznyk, A. V. V.
Nowak, G. *see* Gerlach, K.
Nuño de la Rosa, V. *see* Hernández-Palazón, J.

O'Brien, J. *see* Tyther, R.
O'Rourke, J., Fahy, C. & Donnelly, M. Subcutaneous emphysema at the site of central line placement due to the haematogenous spread of *Clostridium septicum*, 162
Obara, H. *see* Mikawa, K.
Oberthür, A. *see* Hoffmann, E.
Oczenski, W. *see* Fitzgerald, R. D.
Oczenski, W. *see* Krenn, H.
Oehmke, M. J. *see* Knolle, E.
Ogawa-Okamoto, C., Saito, S., Nishihara, F., Yuki, N. & Goto, F. Blood pressure control with glyceryl trinitrate during electroconvulsive therapy in a patient with cerebral aneurysm, 70
Okazaki, Y. *see* Oshima, T.
Okuda, K., Takanishi, T., Yoshimoto, K. & Ueda, S. Trazodone hydrochloride attenuates thermal hyperalgesia in a chronic constriction injury rat model, 409
Okuda, T. *see* Karasawa, F.
Okuda, Y., Urabe, K. & Kitajima, T. Retropharyngeal or cervicomediastinal haematomas following stellate ganglion block, 757
Onimus, T. *see* Lebuffe, G.
Oral, U. *see* Özer, Z.
Oshima, T. *see* Karasawa, F.
Oshima, T., Karasawa, F., Okazaki, Y., Wada, H. & Satoh, T. Effects of sevoflurane on cerebral blood flow and cerebral metabolic rate of oxygen in human beings: a comparison with isoflurane, 543
Otelcioglu, S. *see* Reisli, R.
Özalevli, M. *see* Ünlügenç, H.
Özalp, G. *see* Kaya, M.
Ozcan, S. *see* Apan, A.
Ozcan, S. *see* Başar, H.
Özer, Z., Görür, K., Altunkan, A. A., Bilgin, E., Çamdeviren, H. & Oral, U. Efficacy of tramadol versus meperidine for pain relief and safe recovery after adenotonsillectomy, 920
Ozyar, B. *see* Alper, I.

Ozyar, B. *see* Erhan, E.
Özyilmaz, M. A. *see* Turhanoğlu, S.

Pace, N. *see* Steven, M.
Pallett, E. J. *see* Smart, J. A.
Palmaers, T. *see* Fechner, J.
Paloheimo, M. P. J. *see* Santanen, O. A. P.
Pamukçu, Z. *see* Turan, A.
Panchal, I. D. *see* Shah, F. R.
Papageorgiou, E. P. *see* Mentzelopoulos, S. D.
Paqueron, X. *see* Aubrun, F.
Parfrey, N. *see* Murray, C.
Paris, A. *see* von Knobelsdorff, G.
Parry, M. *see* White, S.
Parsons, R. S. *see* Naughton, C.
Paul, M. *see* Hoffmann, E.
Payne, K., Moore, E. W., Elliott, R. A., Moore, J. K. & McHugh, G. A. Anaesthesia for day case surgery: a survey of paediatric clinical practice in the UK, 325
Payne, K., Moore, E. W., Elliott, R. A., Pollard, B. J. & McHugh, G. A. Anaesthesia for day case surgery: a survey of adult clinical practice in the UK, 311
Peduto, V. A., Baroncini, S., Montanini, S., Proietti, R., Rosignoli, L., Tufano, R. & Casati, A. A prospective, randomized, double-blind comparison of epidural levobupivacaine 0.5% with epidural ropivacaine 0.75% for lower limb procedures, 979
Penzo, D. *see* Guarracino, F.
Perry, E. K. *see* Hanning, C. D.
Peters, J. *see* Lutz, J. T.
Pezza, M. *see* De Robertis, E.
Picker, O., Schwarte, L. A., Schindler, A. W. & Scheeren, T. W. L. Desflurane increases heart rate independent of sympathetic activity in dogs, 945
Picton, P. & Barnes, N. W. More resistance than normal: equipment checks revisited, 503
Piepenbrock, S. *see* Raymondos, K.
Plainer-Zöchling, E. *see* Krenn, H.
Poddar, R. K. & Gosavi, C. P. Analgesic effects of intrathecal neostigmine 984
Pollard, B. J. *see* Payne, K.
Pollard, B. J., Elliott, R. A. & Moore, E. W. Anaesthetic agents in adult day case surgery, 1
Porter, J. M., Markos, F., Snow, H. M. & Shorten, G. D. The efficacy of nicorandil, calcium chloride and nitroglycerin in treatment of ropivacaine-induced cardiotoxicity, 939
Pourriat, J.-L. *see* Kierzek, G.
Powroznyk, A. *see* Naughton, C.
Powroznyk, A. V. V., Vuylsteke, A., Naughton, C., Misso, S. L., Holloway, J., Jolin-Mellgård, Å., Latimer, R. D., Nordlander, M. & Feneck, R. O. Comparison of clevidipine with sodium nitroprusside in the control of blood pressure after coronary artery surgery, 697
Pradier, O. *see* Fattorutto, M.
Proietti, R. *see* Peduto, V. A.
Puig, M. M. *see* Gallart, L.
Puri, G. D. & Murthy, S. S. Bispectral index monitoring in patients undergoing cardiac surgery under cardiopulmonary bypass, 451

Quintard, H. *see* Galinski, M.
Quinzio, L., Junger, A., Gottwald, B., Benson, M., Hartmann, B., Jost, A., Banzhaf, A. & Hempelmann, G. User acceptance of an anaesthesia information management system, 967

Rabey, P. G. *see* Gill, P. S.
Raeder, J. *see* Thagaard, K. S.
Rahman, R. *see* Farmery, A. D.
Rall, M., Mönk, S., Mather, S., Rettedal, A. & Glavin, R. SESAM – The Society in Europe for Simulation Applied to Medicine, 763
Ramírez, G. *see* Hernández-Palazón, J.
Raymondos, K., Münte, S., Krauss, T., Grouven, U. & Piepenbrock, S. Cortical activity assessed by Narcotrend® in relation to haemodynamic responses to tracheal intubation at different stages of cortical suppression and reflex control, 44
Redmond, H. P. *see* Tyther, R.
Reeh, P. W. *see* Blunk, J. A.
Reichle, F. *see* Heindl, B.
Reilly, N. *see* Naughton, C.
Reinecke, H., Breithardt, G. & Van Aken, H. Cardiological aspects in preoperative anaesthesiological evaluation: old heroes, new shadows, 595
Reis, E. *see* de Assunção Braga, A. F.
Reisli, R., Celik, J., Tuncer, S., Yosunkaya, A. & Otelcioglu, S. Anaesthetic and haemodynamic effects of continuous spinal versus continuous epidural anaesthesia with prilocaine, 26
Rellos, K. *see* Mentzelopoulos, S. D.
Rettedal, A. *see* Rall, M.
Reuter, D. A., Felbinger, T. W., Schmidt, C., Moerstedt, K., Kilger, E., Lamm P. & Goetz A. E. Trendelenburg positioning after cardiac surgery: effects on intrathoracic blood volume index and cardiac performance, 17
Riou, B. *see* Aubrun, F.
Robles, R. *see* Hernández-Palazón, J.
Roche, R. J., Farmery, A. D. & Garrard, C. S. Outcome for cardiothoracic surgical patients requiring multidisciplinary intensive care, 719
Roekaerts, P. M. H. J. *see* Heijmans, J. H.
Roepcke, H. *see* Bouillon, T.
Roewer, N. *see* Kranke, P.
Rolf, N. *see* Weber, T. P.
Ronayne, M. *see* Iohom, G.
Rosen, P. *see* Farmery, A. D.
Rosignoli, L. *see* Peduto, V. A.
Rossaint, R. *see* Kuhlen, R.
Rosseel, P. M. J. *see* El Azab, S. R.
Roughan, J. V. *see* Antunes, L. M.
Rozario, C. *see* Harmon, D.
Rubulotta, F. *see* Leykin, Y.
Rutten, C. L. G. *see* van Klei, W. A.

Saager, L. *see* Gerlach, K.
Sade, B. *see* Taplu, A.
Safranski, J. *see* Baigel, G.
Saito, S. *see* Ogawa-Okamoto, C.
Saito, S. *see* Usui, T.
Sakic, K. *see* Kvolik, S.
Sakka, S. G. & Meier-Hellmann, A. Intrathoracic blood volume in a patient with pulmonary embolism, 256
Saklayan, M. *see* Erhan, E.
Sakuragi, T. *see* Harada, C.
Saletu, B. *see* Hellwagner, K.
Salihoglu, Z., Demiroluk, S. & Dikmen, Y. Respiratory mechanics in morbid obese patients with chronic obstructive pulmonary disease and hypertension during pneumoperitoneum, 658
Salmons, V. A. *see* Hollmann, M. W.
Sánchez-Etayo, G., Ayuso, A., Santos, P., Tercero, F. J. & Luis, M. Subcutaneous emphysema, pneumomediastinum and pneumothorax after laryngeal laser surgery, 753
Santanen, O. A. P., Svartling, N., Haasio, J. & Paloheimo, M. P. J. Neural nets and prediction of the recovery rate from neuromuscular block, 87
Santos, P. *see* Sánchez-Etayo, G.
Sariyildiz, O. *see* Kaya, M.

Sato, S. *see* Kobayashi, S.
Satoh, T. *see* Karasawa, F.
Satoh, T. *see* Oshima, T.
Scheeren, T. W. L. *see* Picker, O.
Scheffer, G. J. *see* El Azab, S. R.
Schindler, A. W. *see* Picker, O.
Schmelz, M. *see* Blunk, J. A.
Schmidt, C. *see* Reuter, D. A.
Schmieder, K., Schregel, W., Engelhardt, M., Harders, A. & Cunitz, G. Cerebral vascular reactivity response to anaesthetic induction with propofol in patients with intracranial space-occupying lesions and vascular malformations, 457
Schmitz, A. *see* Gerlach, K.
Schmucker, P. *see* Gerlach, K.
Schneider, G., Elidrissi, C. & Sebel, P. S. Bispectral index-guided administration of anaesthesia: comparison between remifentanil/propofol and remifentanil/isoflurane, 624
Scholz, J. *see* Ledowski, T.
Scholz, J. *see* von Knobelsdorff, G.
Scholz, J. *see* Wappler, F.
Schregel, W. *see* Schmieder, K.
Schroegendorfer, K. *see* Hellwagner, K.
Schulte am Esch, J. *see* von Knobelsdorff, G.
Schulte am Esch, J. *see* Wappler, F.
Schüttler, J. *see* Fechner, J.
Schüttler, J. *see* Schwilden, H.
Schwarte, L. A. *see* Picker, O.
Schweitzer, E. *see* Fitzgerald, R. D.
Schwilden, H., Fechner, J., Albrecht, S., Hering, W., Ihmsen, H. & Schüttler, J. Testing and modelling the interaction of alfentanil and propofol on the EEG, 363
Schwilden, H. *see* Schywalsky, M.
Schywalsky, M., Ihmsen, H., Tzabazis, A., Fechner, J., Burak, E., Vornov, J. & Schwilden, H. Pharmacokinetics and pharmacodynamics of the new propofol prodrug GPI 15715 in rats, 182
Sebel, P. S. *see* Schneider, G.
Seifert, F. *see* Blunk, J. A.
Servillo, G. *see* De Robertis, E.
Shah, F. R., Halbe, A. R., Panchal, I. D. & Goodchild, C. S. Improvement in postoperative pain relief by the addition of midazolam to an intrathecal injection of buprenorphine and bupivacaine, 904
Shannon, P. E. *see* Villevieille, T.
Sharma, P. *see* Singh, B.
Sherry, E. *see* Naughton, C.
Shiga, M. *see* Mikawa, K.
Shlugman, D. *see* Farmery, A. D.
Shorrab, A. A. & Atallah, M. M. Total intravenous anaesthesia with ketamine–midazolam versus halothane–nitrous oxide–oxygen anaesthesia for prolonged abdominal surgery, 925
Shorten, G. D. *see* McDonnell, C. G.
Shorten, G. D. *see* Murray, C.
Shorten, G. D. *see* Porter, J. M.
Shorten, G. *see* Tyther, R.
Simpson, P. Training, assessment and accreditation in anaesthesiology and the implications for the European Union, 679
Singh, B., & Sharma, P. Posterior subgluteal approach to block the sciatic nerve: the controversy of 'which came first, the chicken or the egg' is alive 496
Slappendel, R. *see* Weber, E. W. G.
Smart, J. A., Pallett, E. J. & Duthie, D. J. R. Breath interval as a continuous measure of opioid effects of intravenous fentanyl and alfentanil, 498
Snow, H. M. *see* Murray, C.
Snow, H. M. *see* Porter, J. M.
Spackman, D. *see* Naughton, C.
Spruit, M. *see* Weber, E. W. G.
Stanley, T. Human immobilization: is the experience in Moscow just the beginning?, 427
Steine, S. *see* Thagaard, K. S.
Steven, M. & Pace, N. Non-heart-beating organ donation, 855
Strümper, D. *see* Hollmann, M. W.
Struthers, A. D. *see* Macmillan, C. S. A.
Struys, M. M. R. F. *see* Moerman, A. T.
Strzelczyk, J. *see* Gaszyn´ski, T.
Stuck, B. A. *see* Dahn, J.
Sturgess, J. E., & Leary, T. S. Life-threatening airway obstruction caused by cervical osteophytes, 988
Sungur, Z. *see* Tugrul, M.
Süt, N. *see* Karamanlıoğlu, B.
Süzer, Ö. *see* Bozkurt, P.
Svartling, N. *see* Santanen, O. A. P.
Szmuk, P. *see* Ezri, T.

Takahashi, T. *see* Karasawa, F.
Takamatsu, I. *see* Karasawa, F.
Takanishi, T. *see* Okuda, K.
Takita, A. *see* Karasawa, F.
Tamura, M., Nakamura, K., Kitamura, R., Kitagawa, S., Mori, N. & Ueda, Y. Oral premedication with fentanyl may be a safe and effective alternative to oral midazolam, 482
Taplu, A., Gökmen, N., Erbayraktar, S., Sade, B., Erkan, N., Karadibak, K., Arkan, A. Effects of pressure- and volume-controlled inverse ratio ventilation on haemodynamic variables, intracranial pressure and cerebral perfusion pressure in rabbits: a model of subarachnoid haemorrhage under isoflurane anaesthesia, 690
Tegazzin, V. *see* Wappler, F.
Terbeck, S. *see* Kuhlen, R.
Tercero, F. J. *see* Sánchez-Etayo, G.
Thagaard, K. S., Steine, S. & Raeder, J. Ondansetron disintegrating tablets of 8mg twice a day for 3 days did not reduce the incidence of nausea or vomiting after laparoscopic surgery, 153
Thudium, A. *see* Klockgether-Radke, A. P.
Tok, D. *see* Turhanoğlu, S.
Tomatir, E. *see* Erbay, H.
Tonner, P. H. *see* von Knobelsdorff, G.
Torri, G. *see* Casati, A.
Torri, G. *see* Mascotto, G.
Tortosa, J. A. *see* Hernández-Palazón, J.
Townley, S. A. Central venous catheter malposition in an anomalous pulmonary vein, 985
Tramoni, G., Viale, J. P., Cazals, C. & Bhageerutty, K. Morphine-sparing effect of nefopam by continuous intravenous injection after abdominal surgery by laparotomy, 990
Tran, D. *see* Galinski, M.
Tsutsui, M. *see* Karasawa, F.
Tufano, R. *see* De Robertis, E.
Tufano, R. *see* Peduto, V. A.
Tugrul, M., Camci, E. & Sungur, Z. Effects of sevoflurane and isoflurane on the efficacy of rewarming from hypothermic cardiopulmonary bypass, 583
Tulunay, M. *see* Aşık, I.
Tuncer, S. *see* Reisli, R.
Turan, A. *see* Karamanlıoğlu, B.
Turan, A., Karamanlıoğlu, B., Memiş, D. & Pamukçu, Z. Alternative application site of transdermal nitroglycerin and the reduction of pain on propofol injection, 170
Turhanoğlu, S., Kararmaz, A., Özyilmaz, M. A., Kaya, S. & Tok, D. Effects of different doses of oral ketamine for premedication of children, 56

Turley, A. *see* Campbell, A. E.
Tyagi, A. & Bhattacharya, A. A reply 255
Tyther, R., O'Brien, J., Wang, J., Redmond, H. P. & Shorten, G. Effect of sevoflurane on human neutrophil apoptosis, 111
Tzabazis, A. *see* Schywalsky, M.
Tzanova, I. *see* Wappler, F.
Tzoufi, M. J. *see* Mentzelopoulos, S. D.

Ueda, S. *see* Okuda, K.
Ueda, Y. *see* Tamura, M.
Ugur, G. *see* Alper, I.
Ugur, G. *see* Erhan, E.
Uhlig, T. *see* Gerlach, K.
Unlugenc, H., Guler, T., Gunes, Y., Işik, G. Comparative study of the antiemetic efficacy of ondansetron, propofol and midazolam in the early postoperative period, 668
Ünlügenç, H., Özalevli, M., Güler, T. & Işik, G. Postoperative pain management with intravenous patient-controlled morphine: comparison of the effect of adding magnesium or ketamine, 416
Urabe, K. *see* Okuda, Y.
Urban, B. W. *see* Wartenberg, H. C.
Urwyler, A. *see* Wappler, F.
Usui, T., Saito, S. & Goto, F. Spontaneous intracranial hypotension treated with a cervical epidural blood patch, 500

Vallet, B. *see* Lebuffe, G.
Van Aken, H. *see* Menzebach, A.
Van Aken, H. *see* Reinecke, H.
Van Aken, H. *see* Weber, T. P.
Van Aken, H. *see* Westphal, M.
van der Heide, H. *see* Weber, E. W. G.
van Klei, W. A., Grobbee, D. E., Rutten, C. L. G., Hennis, P. J., Knape, J. T. A., Kalkman, C. J. & Moons, K. G. M. Role of history and physical examination in preoperative evaluation, 612
Van Pelt, F. D. *see* McDonnell, C. G.
Van Praet, F. *see* Cammu, G.
Van Strik, R. *see* El Azab, S. R.
Van Wijk, E. M. *see* El Azab, S. R.
Vardanega, A. *see* Guarracino, F.
Viale, J. P., *see* Tramoni, G.
Vilaplana, J. *see* Arxer, A.
Viljakka, T. J. *see* Haltiavaara, K. M.
Villalonga, A. *see* Arxer, A.
Villevieille, T., Mercier, F., Shannon, P. E., Auroy, Y. & Benhamou, D. Efficacy of epidural analgesia during labour and delivery: a comparison between singleton vertex presentation, singleton breech presentation and twin pregnancies, 164
Vinciguerra, F. *see* Casati, A.
Viscidi, D. *see* De Robertis, E.
von Knobelsdorff, G., Höppner, R. M., Tonner, P. H., Paris, A., Nienaber, C. A., Scholz, J. & Schulte am Esch, J. Induced arterial hypotension for interventional thoracic aortic stent-graft placement: impact on intracranial haemodynamics and cognitive function, 134
Vornov, J. *see* Schywalsky, M.
Vuylsteke, A. *see* Powroznyk, A. V. V.

Wada, H. *see* Karasawa, F.
Wada, H. *see* Oshima, T.
Walsh, K. H., Murphy, C., Iohom, G., Cooney, C. & McAdoo, J. Comparison of the effects of two intrathecal anaesthetic techniques for transurethral prostatectomy on haemodynamic and pulmonary function, 560
Wang, J. *see* Tyther, R.
Wappler, F., Anetseder, M., Baur, C. P., Censier, K., Doetsch, S., Felleiter, P., Fiege, M., Fricker, R., Halsall, P. J., Hartung, E., Heffron, J. J. A., Heytens, L., Hopkins, P. M., Klingler, W., Lehmann-Horn, F., Nivoche, Y., Tegazzin, V., Tzanova, I., Urwyler, A., Weißhorn, R., Schulte am Esch, J. Multicentre evaluation of in vitro contracture testing with bolus administration of 4-chloro-*m*-cresol for diagnosis of malignant hyperthermia susceptibility, 528
Wappler, F., Frings, D. P., Scholz, J., Mann, V., Koch, C. & Schulte am Esch, J. Inhalational induction of anaesthesia with 8% sevoflurane in children: conditions for endotracheal intubation and side-effects, 548
Wartenberg, H. C., Wartenberg, J. P. & Urban, B. W. Pharmacological modification of sodium channels from the human heart atrium in planar lipid bilayers: electrophysiological characterization of responses to batrachotoxin and pentobarbital, 354
Wartenberg, J. P. *see* Wartenberg, H. C.
Warters, R. D. *see* Ezri, T.
Waschke, K. F. *see* Dahn, J.
Washington, J. M. *see* Hollmann, M. W.
Weber, E. W. G., Slappendel, R., Durieux, M. E., Dirksen, R., van der Heide, H. & Spruit, M. COX 2 selectivity of non-steroidal anti-inflammatory drugs and perioperative blood loss in hip surgery. A randomized comparison of indomethacin and meloxicam, 963
Weber, T. P., Große Hartlage, M. A., Van Aken, H. & Booke, M. Anaesthetic strategies to reduce perioperative blood loss in paediatric surgery, 175
Weber, T. P., Van Aken, H., Kehrel, B. E., Meißner, A., Brüssel, T., Bullmann, V., Winkelmann, W., Heindel, W. & Rolf, N. Epidural bleed and quadriplegia due to acquired platelet dysfunction unrelated to multiple spinal and epidural puncture, 333
Wedley, J. R. *see* Ather, M.
Weindlmayr-Goettel, M. *see* Hellwagner, K.
Weindlmayr-Goettel, M. *see* Ledowski, T.
Weißhorn, R. *see* Wappler, F.
Westphal, M., Hohage, H., Buerkle, H., Van Aken, H., Ermert, T. & Brodner, G. Adsorption of sufentanil to epidural filters and catheters, 124
Whitaker, D. C. Acid–base management during hypothermic neuroprotection, 422
White, M. *see* Heffernan, A. M.
White, S., Parry, M. & Henderson, K. Anaesthesia for total hip replacement in a patient with Holt–Oram syndrome, 336
Wielogorski, A. *see* Naughton, C.
Wilkes, A. R. *see* Campbell, A. E.
Wilkes, M. M. *see* Haynes, G. R.
Winkelmann, W. *see* Weber, T. P.
Wouters, P. F. *see* Leather, H. A.
Wulf, H. *see* Ledowski, T.

Yamada, S. *see* Karasawa, F.
Yanagidate, F. & Dohi, S. Two episodes of bronchospasm during anaesthesia in asthmatic patients, 579
Yegin, A., Yilmaz, M., Karsli, B. & Erman, M. Analgesic effects of intrathecal neostigmine in perianal surgery, 404
Yelken, B. *see* Bilir, L.
Yilmaz, M. *see* Yegin, A.
Yörükoğlu, D. *see* Aşık, I.
Yoshimoto, K. *see* Okuda, K.
Yosunkaya, A. *see* Reisli, R.
Yüceyar, L. *see* Erolçay, H.
Yuki, N. *see* Ogawa-Okamoto, C.

Ziade, M. F. *see* Naja, M. Z.
Zuokumor, P. & Columb, M. Epidural anaesthesia for pain relief in labour, 674